

## Short communication

## Paleosols and pedostratigraphy

Edoardo A.C. Costantini

*Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria, Research Centre for Agriculture and Environment, Florence, Italy*

## ARTICLE INFO

## Keywords:

Definition  
Palaeo  
Soil  
Geosol  
Method  
Stratigraphy  
UBSU  
Scopus  
Cultural heritage  
Red list  
Climate change  
Pedosites  
Biodiversity

## ABSTRACT

Paleopedology and pedostratigraphy are devoted to the use of buried and relict soils in interpreting the record of environmental conditions that are different from the current ones. As such, paleosols may be used in models that aim at predicting future earth system reactions to changing environmental conditions. Dated paleosols are an important part of the natural heritage and should be protected.

## 1. Paleosols

Paleosols are soils that formed, or started to form, in times earlier than recent and often under environmental conditions that were different from the current ones. They show horizons and features diverse from those of younger soils, formed in analogous present settings, because of a change of at least one soil forming factor, lasting enough time and acting so strong to affect the kind or rate of pedogenesis. Actually, many paleosols began to develop long periods ago and have little relationship in their chemical and physical characteristics to the present-day climate, vegetation, or topographic position.

We distinguish relict and buried paleosols (Retallack, 1998). Relict paleosols remain at the surface today and continue to develop, while buried paleosols have been buried by younger sediments such as loess, glacial till, colluvium or alluvium, and can be more or less modified from the burial and even diagenesed or petrified (Olson and Nettleton, 1998). Burial can cause modification of soil properties and features, especially in the uppermost horizons. An intermediate possibility is that of welded paleosols. Welded paleosols show sediments that have been pedogenetically incorporated in the soil profile, since the covering layer (e.g. aeolian, fluvial, colluvial) is not thick enough to isolate the underlying soil profile from pedogenesis. Such profiles can be called “Polygenetic”.

Paleosols are a fundamental source of evidence for environmental processes that took place in the past and, as such, they may be used in

models that aim at predicting future earth system reactions to changing environmental conditions. Large numbers of papers on paleopedology are continuously being published. We observe a progressive increase in the number of papers during the last 30 years (Fig. 1), especially in the last years, driven by the concern about climate change and the interest on the reaction of the earth system during analogue past climate changes (Sauer et al., 2008a,b), but also by the need to steer their management. In fact, paleosols frequently have low fertility and some limiting condition to agricultural use, because of long and deep weathering and leaching of nutrient (Retallack, 1978). However, moderate fertility of paleosols with low-grade plinthite has been found functional to produce high quality wine (Costantini and Priori, 2007) and some of the best terroir correspond to paleosols (Costantini et al., 2012; Retallack and Burns, 2016). In addition, where paleosols do not dominate the landscape, likewise in countries without a tropical climate, they can host endemisms, which increase the biodiversity of a land. Actually, an empirical correlation between paleosol diffusion, pedodiversity, and biodiversity, has been demonstrated (Ibañez et al., 1995; Costantini and L'Abate, 2016a,b).

Along with paleosols genesis, the focuses of paleopedological research are:

- i) the reconstruction of paleo environmental characteristics and development, in particularly climate. That is, the use of soil characteristics as proxies of palaeo-environmental conditions and time

E-mail address: [edoardo.costantini@crea.gov.it](mailto:edoardo.costantini@crea.gov.it).

<http://dx.doi.org/10.1016/j.apsoil.2017.09.021>

Received 21 December 2016; Received in revised form 12 September 2017; Accepted 14 September 2017

Available online 04 November 2017

0929-1393/ © 2017 Elsevier B.V. All rights reserved.




Fig. 1. Papers dealing with paleosols during the years 1986–2015 (Scopus database). Published works and cumulative curve.

of surface formation (a.o. Costantini and Damiani, 2004; Sauer, 2010; Carnicelli and Costantini, 2013);

- ii) the quantification of pedogenic processes, studied in soil chronosequences established on sequences of river terraces (a.o. Torrent, 1976; Arduino et al., 1984; Ajmone Marsan et al., 1988; Bain et al., 1993; Tsai et al., 2006), marine terraces (a.o. Muhs, 1982; Scarciglia et al., 2006; Tsai et al., 2007; Wagner et al., 2007; Sauer et al., 2010), beach ridges (a.o. Sauer et al., 2007; Sauer et al., 2008a,b), moraines (a.o. Egli et al., 2001), dunes of different ages (a.o. Lichter, 1998), volcanic (a.o., Dazzi, 2007; Jahn et al., 1987; Solleiro-Rebolledo et al., 2015), colluvial (a.o. Costantini et al., 2006), or loess depositions (a.o. Bronger, 2003; Costantini et al., 2009; Sauer et al., 2015);
- iii) Heritage soil documentation and protection. Paleosols store information about the environmental conditions during their genesis and thus reflect the natural and cultural heritage of the landscape. This particular nature of paleosols, referred to as “soil memory” (Targulian and Goryachkin, 2004; Costantini et al., 2007a), makes them particularly valuable among the various earth soil bodies.

## 2. Pedostratigraphy

Adding soil and paleosol information to geological mapping has always been a challenge for both paleopedologists and geologists (Morrison, 1993). The North American Commission on Stratigraphic Nomenclature (1983) has defined the concept and application of pedostratigraphic unit as “a buried, traceable, three-dimensional body of rock that consists of one or more differentiated pedologic horizons” (UBSU. Unconformity-bounded Stratigraphic Units). The upper and lower boundaries of a pedostratigraphic unit correspond to the soil horizons formed by pedogenesis in a buried soil profile. A pedostratigraphic unit is characterized by the range of physical and chemical properties in the type area, rather than by “typical” properties exhibited in a type section. A pedostratigraphic unit has a consistent stratigraphic position and is defined at a type locality where the horizons are buried by younger deposits, but may be traced to sites where it crops out on the present land surface. The term “pedostratigraphic unit” is frequently considered a synonym of “Geosol” (WG on Definitions used in Paleopedology, 1997). Using the Geosol concept, inter-regional or even global correlation of buried paleosols, and loess-paleosol sequences, were proposed by Rutter et al. (1991), Benvenuti et al. (2002), Bronger (2003), Jacobs et al. (2009), Marković et al. (2015), Sycheva and Khokhlova (2016).

Costantini et al. (2007b) worked out the concept of “pedostratigraphic level”, which differentiates from that of “pedostratigraphic

unit” or “Geosol” since it can be applied to non-buried paleosols. The rationale is that the whole soil body, from the soil surface downwards, can be classified in levels according to genetic characteristics and degree of weathering. Therefore, a pedostratigraphic level (PL) is considered as a characteristic assemblage of similar genetic soil horizons, formed by materials having the same degree of weathering, and an estimated age through correlation to benchmark soils. Thus a PL is a three-dimensional soil layer with:

1. traceable upper and lower boundaries
2. showing a characteristic assemblage of similar soil genetic horizons (e.g., Bt1, Bt2, etc.)
3. formed in materials having a certain degree of weathering (e.g., clay minerals assemblage, free crystalline iron content) and
4. having an age that is estimated by means of direct dating or correlation with dated horizons.

This approach needs the characterization of soil profile horizons according to two criteria: i) the expression of pedofeatures, leading to the identification of a class of genetic horizon, and ii) the degree of weathering of the parent material. Both criteria are related to the five soil forming factors, but the second one provides better clues on the duration of pedogenesis. In addition, the decoupling of the two criteria allows to clearly separate pedofeatures formed either in fresh or pre-weathered materials. This can be useful to distinguish, for instance, soils formed from unweathered sediments deposited in-situ and not reworked, from those developed in colluvia, where already weathered soil sediments accumulated.

Similar to mapping Geosols in buried paleosols, tracing pedostratigraphic levels (PLs) in relict paleosols is a four-dimensional approach in the study of soils and associated sediments, integrating field and laboratory analyses. It takes into account both horizontal as well as vertical field soil variations and time pegs determined in the laboratory.

The methodology can be summarized in the following 8 steps:

1. soils are placed in their geomorphological context,
2. sedimentological processes are recognized on natural or artificial sections,
3. soil profiles are described and analysed, in order to identify genetic processes characterizing each soil horizon and to highlight discontinuities,
4. the degree of weathering of the parent material of each soil horizon is assessed, by means of selected aging indicators,
5. time pegs are fitted by means of laboratory datings and proxies,
6. horizons with same genetic processes and degree of weathering of




Fig. 2. A paleosols chronostratigraphy on a marine terrace sequence in southern Sicily (from Wagner et al., 2007). The soils of the catena differentiate for several features, according to their age and local setting. Under the Mediterranean climate, colours below the darker ploughed layer tend to become redder first, and then yellowish or blackish, along with the accumulation of iron, manganese, and other metals in different forms (Costantini et al., 2006). In parallel, soil depth increases up to several meters, through the weathering of the original parent materials, that is, calcarenite and marine limestone of different geological eras, with some covers of aeolian and colluvial sediments.

the parent material are assembled in progressive PLs,

7. each PL is correlated with dated horizon of benchmark profiles, showing same genetic properties and degree of weathering of the parent material,
8. PLs belonging to different soil profiles are correlated (Fig. 2).

### 3. Conclusions

The large wealth of papers documenting occurrence and extent of paleosols all over the world demonstrates that this kind of soils are more common than usually reported on soil maps. This fact has several practical consequences, in particular, on soil mapping and interpretation at the detailed scales, since paleosols features and their relationship with present soil forming factors do not follow the pattern of modern soils. For instance, the reliability of inference systems to digitally map the soil cover using as predictive layers current morphology, vegetation, and climate, may be very poor. In addition, algorithms that model soil features along the profile may also provide inconsistent results, because of the presence of several genetic and lithological discontinuities.

The pedomorphological approach has been successfully used for detailed reconstructions, applied to various environments and for different scopes. However, there are also weak points. Specifically, it is more suitable for local than for regional correlations, it needs costly laboratory analyses and specific expertise, and it gives mainly indirect chronostratigraphy. In the next future, it is foreseeable that pedostratigraphy and paleopedological studies will be boosted by improvements in dating techniques and proxies calibration.

Dated paleosols are considered particularly important, in that they can be used as stratigraphic markers and thus give more accuracy to the mapping of Quaternary formations. Dated paleosols, likewise soils from

the archeological and paleontological sites, are an important part of the natural heritage and should be protected (Costantini and L'Abate, 2009).

In Italy, most relevant pedosites (that is, cultural heritage sites of soil nature) can be overlaid on Google Earth, by downloading the file "pedosites\_it.kml" from [http://www.soilmaps.it/download/dt-Soilsites\\_of\\_Italy\\_26.kml](http://www.soilmaps.it/download/dt-Soilsites_of_Italy_26.kml). It seems possible and indeed desirable that similar repositories of pedosites will be produced in other European countries, especially where paleosols possessing cultural heritage are threatened by urbanization, to complete a red list of the European soils having cultural heritage and in danger of extinction and, in perspective, to extend it at the global scale.

### References

- Ajmoné Marsan, F., Barberis, E., Arduino, E., 1988. A soil chronosequence in north-western Italy: morphological, physical and chemical characteristics. *Geoderma* 42, 51–64.
- Arduino, E., Barberis, E., Carraro, F., Forno, M.G., 1984. Estimating relative ages from iron-oxide/total-iron ratios of soils in the Western Po Valley, Italy. *Geoderma* 33, 39–52.
- Bain, D.C., Mellor, A., Robertson-Rintoul, M.S.E., Buckland, S.T., 1993. Variations in weathering processes and rates with time in a chronosequence of soils from Glen Feshie, Scotland. *Geoderma* 57, 275–293.
- Benvenuti, 2002 Benvenuti, M., Carnicelli, S., Belluomini, G., Dainelli, N., Di Grazia, S., Ferrari, G.A., Iasio, C., Sagr, M., Ventra, D., Anafi, B., 2002. The Ziway-Shala lake basin (main Ethiopian rift, Ethiopia): a revision of basin evolution with special reference to the Late Quaternary. *J. Afr. Earth. Sci.* 2, 247–269.
- Bronger, A., 2003. Correlation of loess-paleosol sequences in East and Central Asia with SE Central Europe: towards a continental Quaternary pedomorphology and paleoclimatic history. *Quat. Int.* 106–107, 11–31.
- Carnicelli, S., Costantini, E.A.C., 2013. Time as a soil forming factor and age of Italian soils. In: Costantini, E.A.C., Dazzi, C. (Eds.), *The Soils of Italy*. World Soils Book Series. Springer, pp. 93–104.
- Costantini, E.A.C., Damiani, D., 2004. Clay minerals and the development of Quaternary soils in central Italy. *Rev. Mex. Cienc. Geol.* 21 (1), 144–159.

- Costantini, E.A.C., L'Abate, G., 2009. The soil cultural heritage of Italy: geodatabase maps, and pedodiversity evaluation. *Quat. Int.* 209, 142–153.
- Costantini, E.A.C., L'Abate, G., 2016a. Beyond the concept of dominant soil: preserving pedodiversity in upscaling soil maps. *Geoderma* 271 (1 June 2016), 243–253. <http://dx.doi.org/10.1016/j.geoderma.2015.11.024>.
- Costantini, E.A.C., L'Abate, G., 2016b. Beyond the concept of dominant soil: preserving pedodiversity in upscaling soil maps. *Geoderma* 271 (1 June 2016), 243–253. <http://dx.doi.org/10.1016/j.geoderma.2015.11.024>.
- Costantini, E.A.C., Priori, S., 2007. Pedogenesis of plinthite during early Pliocene in the Mediterranean environment. Case study of a buried paleosol at Podere Renieri, central Italy. *Catena* 71, 425–443.
- Costantini, E.A.C., Lessovaia, S., Vodyanitskii, Y., 2006. Using the analysis of iron and iron oxides in paleosols (TEM, geochemistry and iron forms) for the assessment of present and past pedogenesis. *Quat. Int.* 156–157, 200–211.
- Costantini, E.A.C., Napoli, R., D'Egidio, G., 2007a. Adding information about soils and paleosols to geological maps, through the application of the “pedostratigraphic level” concept. *Quat. Int.* 175, 125–139.
- Costantini, E.A.C., Malucelli, F., Brenna, S., Rocca, A., 2007b. Using existing soil databases to consider paleosols in land planning. The case study of the Lombardy Region (Northern Italy). *Quat. Int.* 162–163, 166–171.
- Costantini, E.A.C., Priori, S., Urban, B., Hilgers, A., Sauer, D., Protano, G., Trombino, L., Hülle, D., Nannoni, F., 2009. Multidisciplinary characterization of the middle Holocene eolian deposits of the Elsa river basin (central Italy). *Quat. Int.* 209 (1–2), 107–130.
- Costantini, E.A.C., Buccelli, P., Priori, S., 2012. Quaternary landscape history determines the soil functional characters of terroir. *Quat. Int.* 265, 63–73.
- Dazzi, C., 2007. Environmental features and land use of Etna (Sicily-Italy). In: Arnalds, Ö., Bartoli, F., Buurman, P., Öskarsson, H., Stoops, G., García-Rodeja, E. (Eds.), *Soils of Volcanic Regions in Europe*. Springer, pp. 629–644. [http://dx.doi.org/10.1007/978-3-540-48711-1\\_44](http://dx.doi.org/10.1007/978-3-540-48711-1_44). (ISBN 10 3-540-48710-7).
- Egli, M., Fitze, P., Mirabella, A., 2001. Weathering and evolution of soils formed on granitic, glacial deposits: results from chronosequences of Swiss alpine environments. *Catena* 45, 19–47.
- Ibañez, J.J., De-Albs, S., Bermúdez, F.F., García-Álvarez, A., 1995. Pedodiversity: concepts and measures. *Catena* 24 (3), 215–232.
- Jacobs, P.M., Konen, M.E., Curry, B.B., 2009. Pedogenesis of a catena of the Farmdale–Sangamon Geosol complex in the north central United States. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 282 (1), 119–132.
- Jahn, R., Zarei, M., Stahr, K., 1987. Formation of clay minerals in soils developed from basic volcanic rocks under semiarid climatic conditions in Lanzarote, Spain. *Catena* 14 (4), 359–368.
- Lichter, J., 1998. Rates of weathering and chemical depletion in soils across a chronosequence of Lake Michigan sand dunes. *Geoderma* 85, 255–282.
- Marković, 2015 Marković, S.B., Stevens, T., Kukla, G.J., Hambach, U., Fitzsimmons, K.E., Gibbard, P., Buggle, B., Zech, M., Guo, Z., Hao, Q., Wu, H., O'Hara Dhand, K., Smalley, I.J., Újvári, G., Sümege, P., Timar-Gabor, A., Veres, D., Sirocko, F., Vasiljević, D.A., Jary, Z., Svensson, A., Jović, V., Lehmkuhl, F., Kovács, J., Svirčev, A., 2015. Danube loess stratigraphy towards a pan-European loess stratigraphic model. *Earth Sci. Rev.* 148, 228–258.
- Morrison, R.B., 1993. How Can the Treatment of Pedostratigraphic Units in the North American Stratigraphic Code Be Improved? [Online] [www.fadr.msu.ru/inqua/nl-](http://www.fadr.msu.ru/inqua/nl-archive/nl-8-11/toc.html) archive/nl-8-11/toc.html (verified August 2017).
- Muhs, D., 1982. A soil chronosequence on quaternary marine terraces San Clemente Island, California. *Geoderma* 28, 257–283.
- North American Commission on Stratigraphic Nomenclature, 1983. North american stratigraphic code. *Am. Assoc. Petrol. Geol. Bull.* 67 (5), 841–875.
- Olson, C.G., Nettleton, W.D., 1998. Paleosols and the effects of alteration. *Quat. Int.* 51, 185–194.
- Retallack, G.J., Burns, S.F., 2016. The effects of soil on the taste of wine. *GSA Today* 26 (5).
- Retallack, G.J., 1978. Weathering, soils, and paleosols. *Sedimentology*. Springer, Berlin, Heidelberg, pp. 1271–1277.
- Retallack, G.J., 1998. Core concepts of paleopedology. *Quat. Int.* 51, 203–212.
- Rutter, N., Zhongli, D., Evans, M.E., Tungsheng, L., 1991. Baoji-type pedostratigraphic section, loess plateau, north-central China. *Quat. Sci. Rev.* 10 (1), 1–22.
- Sauer, D., Schellmann, G., Stahr, K., 2007. A soil chronosequence in the semi-arid environment of Patagonia (Argentina). *Catena* 71, 382–393.
- Sauer, D., Costantini, E.A.C., Jahn, R., Makeev, A., Feger, K.-H., Felix-Henningsen, P., 2008a. Imprint of environmental change on paleosols. *J. Plant Nutr. Soil Sci.* 171 (4 (August)), 482.
- Sauer, D., Schulli-Maurer, I., Sperstad, R., Sorensen, R., Stahr, K., 2008b. Podzol development with time in sandy beach deposits in southern Norway. *J. Plant Nutr. Soil Sci.* 171 (4), 483–497.
- Sauer, D., Wagner, S., Bruckner, H., Scarciglia, F., Mastronuzzi, G., Stahr, K., 2010. Soil development on marine terraces near Metaponto (Gulf of Taranto, southern Italy). *Quat. Int.* 222 (1–2), 48–63.
- Sauer, D., Amit, R., Carnicelli, S., Costantini, E.A.C., 2015. Rates of soil forming processes and the role of aeolian influx. *Quat. Int.* 376, 1–4.
- Sauer, D., 2010. Approaches to quantify progressive soil development with time in Mediterranean climate-I. Use of field criteria. *J. Plant Nutr. Soil Sci.* 173 (6), 822–842.
- Scarciglia, F., Pulice, I., Robustelli, G., Vecchio, G., 2006. Soil chronosequences on Quaternary marine terraces along the northwestern coast of Calabria (Southern Italy). *Quat. Int.* 156/157, 133–155.
- Solleiro-Rebolledo, E., Sedov, S., Cabadas-Báez, H., 2015. Use of soils and palaeosols on volcanic materials to establish the duration of soil formation at different chronological scales. *Quat. Int.* 376, 5–18.
- Sycheva, S., Khokhlova, O., 2016. Genesis 14C age, and duration of development of the Bryansk paleosol on the Central Russian Upland based on dating of different materials. *Quat. Int.* 399, 111–121.
- Targulian, V.O., Goryachkin, S.V., 2004. Soil memory: types of record, carriers, hierarchy and diversity. *Rev. Mex. Cienc. Geol.* 21, 1–9.
- Torrent, J., 1976. Soil development in a sequence of river terraces in northern Spain. *Catena* 3, 137–151.
- Tsai, H., Huang, W.S., Hseu, Z.Y., Chen, Z.S., 2006. A river terrace soil chronosequence of the Pakua Tableland in central Taiwan. *Soil Sci.* 171, 167–179.
- Tsai, C.C., Tsai, H., Hseu, Z.Y., Chen, Z.S., 2007. Soil genesis along a chronosequence on marine terraces in eastern Taiwan. *Catena* 71, 394–405.
- WG on Definitions used in Paleopedology, 1997. *Paleopedology Glossary*. [Online] <http://fadr.msu.ru/inqua/nl-14/glossary.html> (verified August 2017).
- Wagner, S., Costantini, E.A.C., Sauer, D., Stahr, K., 2007. Soil genesis with time in a marine terrace sequence of Sicily, Italy. *Rev. Mex. Cienc. Geol.* 24, 247–260.